

Asymmetric synthesis of enantiomerically enriched (*S*)- α -aminopropionic acids containing more than one heteroatom in the side-chain radical


A.S. Saghyan^{a,b}, A.S. Dadayan^{a,b}, S.G. Petrosyan^{a,b}, H.M. Simonyan^{a,b},
A.V. Geolchanyan^{a,b}, H.R. Zurnachyan^a, A.F. Mkrtchyan^{a,b}, G.M. Mkrtchyan^b
^aYerevan State University, 1 Alex Manoogian Str, Yerevan, 0025, Armenia
^bSPC "Armbiotechnology" NAN RA, 14 Gyurjyan Str., Yerevan, 0056, Armenia
E-mail: saty_petrosyan@yahoo.com

The classes of heterocyclic compounds known as imidazole, thiazole, thiazole are found in many natural and synthetic products with a wide range of biological activities, and they can be well illustrated by a large number of drugs in the market containing this nucleus [1,2,3]. On the other hand, enantiomerically enriched α - and β -substituted α -amino acids are found in many physiologically active peptides, antibiotics and other drugs[4].

Proceeding from the aforesaid, it is evident that introduction of more than one different heteroatoms containing heterocyclic radicals into the side-chain of amino acids can result in new interesting potentially biologically active compounds.

In this paper we report the asymmetric synthesis of novel β -heterocycle substituted analogs of (*S*)- α -aminopropionic acid containing imidazole, triazole, thiodiazole, thiazole and oxazole substituents in the side-chain radical.

Asymmetric addition of the heterocyclic nucleophiles to the C=C bond of dehydroalanine moiety in Ni^{II} complexes of Schiff's base with chiral auxiliaries (*S*)-2-N-(N'-benzylpropyl)aminobenzophenone proceeded in MeCN in the presence of K₂CO₃ at 20-25°C (Scheme).



As a result diastereomeric mixtures of addition products with a large excess of (*S,S*)-diastereomeric complexes containing amino acids of (*S*)-absolute configuration were formed.

Decomposition of diastereoisomeric complexes was carried out with aqueous HCl, followed by the recovery of the insoluble hydrochlorides of the auxiliary (90%) and purification of the amino acids by ion-exchange method. The target amino acids were isolated with high enantiomeric purity (ee >97%) after crystallization from an aqueous-ethanol (1:2) solution.

1. M.V.N. de Souza // J. Sulfur Chem., 26, 429 (2005).
2. B. Tozcoparan et al. // Bioorganic & Medicinal Chemistry, vol. 15, no. 4, pp. 1808–1814, 2007.
3. J. R. Kumar // Pharmacophore 2010, Vol. 1 (3), 167-177
4. G.C. Barrett, "Chemistry and Biochemistry of the Amino Acids", L.-N.Y.,1985, p.246.

The work has been carried out in the frame of Volkswagenstiftung Project-ID Az 86 223